



# HIGHER ORDER DERIVATIVE-FREE ITERATIVE METHODS WITH AND WITHOUT MEMORY IN BANACH SPACE UNDER WEAK CONDITIONS

Ioannis K. Argyros\* and Santhosh George\*\*

Department of Mathematical Sciences  
Cameron University  
Lawton, OK 73505, US  
E-mails: [iargyros@cameron.edu](mailto:iargyros@cameron.edu)

\*\* Department of Mathematical and Computational Sciences  
NIT Karnataka  
India-575 025  
E-mails: [sgeorge@nitk.ac.in](mailto:sgeorge@nitk.ac.in)

\*Corresponding author.

**Abstract** We study the method considered in Ahmad et al. (2016), for solving systems of nonlinear equations, modified suitably to include the nonlinear equations in Banach spaces. We use the idea of restricted convergence domains instead of Taylor's expansion in our convergence analysis and our conditions are weaker than the conditions used in earlier studies. Numerical examples where earlier results cannot apply to solve equations but our results can apply are also given in this study.

**MSC:** 65J20, 49M15, 74G20, 41A25.

**Keywords:** Newton's method, Divided difference, radius of convergence, local convergence, derivative free method.

---

Submission date: 28 December 2016 / Acceptance date: 16 July 2017 / Available online: 1 August 2017  
Copyright 2017 © Theoretical and Computational Science 2017.

---

## 1. INTRODUCTION

In [8], Ahmad et al. considered the following method

$$\begin{aligned}y_n &= x_n - q_1, \\z_n &= y_n - a_0q_2 - (3 - 2a_0)q_3 - (a_0 - 2)q_4, \\x_{n+1} &= z_n - a_1q_5 - a_2q_6 - a_3q_7 - a_4q_8 - a_5q_9;\end{aligned}\tag{1.1}$$

$$u_n = x_n + b_0F(x_n), A_n = [x_n, u_n, F], h_n = y_n + b_1F(y_n), B_n = [h_n, y_n : F], l_n = z_n + b_2F(z_n), C_n = [l_n, z_n; F], b_0, b_1, b_2 \in \mathbb{R}, A_nq_1 = F(x_n), A_nq_2 = F(y_n), A_nq_3 =$$

---

© 2015 By TaCS Center, All rights reserved.



Published by Theoretical and Computational Science Center (TaCS),  
King Mongkut's University of Technology Thonburi (KMUTT)

$B_n q_2, A_n q_4 = B_n q_3, A_n q_5 = F(z_n), A_n q_6 = C_n q_5, A_n q_7 = C_n q_6, A_n q_8 = C_n q_7, A_n q_9 = C_n q_8, a_1 = 4 + a_5, a_2 = -6 - 4a_5, a_3 = 4 + 6a_5, a_4 = -1 - 4a_5$ . and  $[\cdot, \cdot; F] : D^2 \rightarrow L(X)$  is a divided difference of order one on  $D^2$  [2], for solving systems of equations, provided that  $X = \mathbb{R}^j$  and  $D \subset X$ . In this study we consider method (1.1), but for approximating a locally unique solution  $x^*$  of the nonlinear equation

$$F(x) = 0, \quad (1.2)$$

where  $F$  is a continuous operator defined on a convex subset  $D$  of a Banach space  $X$  with values in  $X$ . The divided difference of order one satisfies  $[x, y; F](x - y) = F(x) - F(y)$  for  $x \neq y$  with  $x, y \in D$  and  $[x, x; F] = F'(x)$ ,  $x \in D$ , if  $F$  is Fréchet-differentiable on  $D$ . We use the notation  $U(u, \rho), \bar{U}(u, \rho)$  to denote the open and closed balls in  $X$ , respectively with center  $u \in X$  and of radius  $\rho > 0$ .

Finding solution for (1.2) is an important problem in mathematics due to the wide application of the equation (1.2). Convergence analysis of the method in [2] used assumptions on the eighth Fréchet-derivative although no derivative appears in the method. This assumption on the higher order Fréchet derivatives of the operator  $F$  restricts the applicability of method (1.1). For example consider the following;

**EXAMPLE 1.1.** Let  $X = C[0, 1], D = \bar{U}(x^*, 1)$  and consider the nonlinear integral equation of the mixed Hammerstein-type [4, 7, 20] defined by

$$x(s) = \int_0^1 G(s, t) \frac{x(t)^2}{2} dt,$$

where the kernel  $G$  is the Green's function defined on the interval  $[0, 1] \times [0, 1]$  by

$$G(s, t) = \begin{cases} (1-s)t, & t \leq s \\ s(1-t), & s \leq t. \end{cases}$$

The solution  $x^*(s) = 0$  is the same as the solution of equation (1.2), where  $F : C[0, 1] \rightarrow C[0, 1]$  is defined by

$$F(x)(s) = x(s) - \int_0^1 G(s, t) \frac{x(t)^2}{2} dt.$$

Notice that

$$\left\| \int_0^1 G(s, t) dt \right\| \leq \frac{1}{8}.$$

Then, we have that

$$F'(x)y(s) = y(s) - \int_0^1 G(s, t)x(t)dt,$$

so since  $F'(x^*(s)) = I$ ,

$$\|F'(x^*)^{-1}(F'(x) - F'(y))\| \leq \frac{1}{8} \|x - y\|.$$

One can see that the work in [2] cannot be applied in this setting, if we choose  $[x, y; F] = \int_0^1 F'(y + \theta(x - y))d\theta$ .

Our goal is to weaken the assumptions in [2] and apply the method for solving equation (1.2) in Banach spaces, so that the applicability of the method (1.1) can be extended. The technique of restricted convergence domains can apply on other iterative methods [1, 3-25].

The rest of the paper is organized as follows. In Section 2 we present the local convergence analysis. We also provide a radius of convergence, computable error bounds and

uniqueness result not given in [2]. Special cases and numerical examples are given in the last section.

## 2. LOCAL CONVERGENCE ANALYSIS

Let  $b_0, b_1, b_2 \in \mathbb{R}$ . Let  $w_0 : [0, +\infty) \times [0, +\infty) \rightarrow [0, +\infty)$ ,  $v_0 : [0, +\infty) \rightarrow [0, +\infty)$  be a continuous, non-decreasing in both variables satisfying  $w_0(0, 0) = 0$ . Define the parameter  $r_0$  by

$$r_0 = \sup\{t \geq 0 : w_0(t, (1 + |b_0|v_0(t))t) < 1\}. \quad (2.1)$$

Let also  $v, w : [0, r_0) \times [0, r_0) \rightarrow [0, +\infty)$  be continuous and non-decreasing functions with  $w(0, 0) = 0$ . Define functions  $g_i, \bar{g}_i$ ,  $i = 1, 2, 3$  on the interval  $[0, r_0)$  by

$$\begin{aligned} g_1(t) &= \frac{w(t, t + |b_0|v_0(t)t)}{1 - p(t)}, \quad p(t) = w_0(t, t + |b_0|v_0(t)t), \\ \bar{g}_1(t) &= g_1(t) - 1, \\ g_2(t) &= \left[ \frac{w(t + g_1(t)t, t + |b_0|v_0(t)t)}{1 - p(t)} + \frac{|1 - a_0|v(g_1(t)t, 0)}{1 - p(t)} \right. \\ &\quad + |3 - 2a_0| \frac{v(g_1(t)t + |b_1|v_0(g_1(t)t)g_1(t)t, g_1(t)tv(g_1(t)t, 0))}{(1 - p(t))^2} \\ &\quad \left. + |a_0 - 2| \frac{v(g_1(t)t + |b_1|v_0(g_1(t)t)g_1(t)t, g_1(t)t^2v(g_1(t)t, 0))}{(1 - p(t))^3} \right] g_1(t), \\ \bar{g}_2(t) &= g_2(t) - 1, \\ g_3(t) &= \left[ \frac{w(t + g_2(t)t, t + |b_0|v_0(t)t)}{1 - p(t)} + \frac{|1 - a_1|v(g_2(t)t, 0)}{1 - p(t)} \right. \\ &\quad + |a_2| \frac{v(g_2(t)t + |b_2|v_0(g_2(t)t)g_2(t)t, g_2(t)t)}{(1 - p(t))^2} \\ &\quad + |a_3| \frac{v(g_2(t)t + |b_2|v_0(g_2(t)t)g_2(t)t, g_2(t)t^2)}{(1 - p(t))^3} \\ &\quad + |a_4| \frac{v(g_2(t)t + |b_2|v_0(g_2(t)t)g_2(t)t, g_2(t)t^3)}{(1 - p(t))^4} \\ &\quad \left. + |a_5| \frac{v(g_2(t)t + |b_2|v_0(g_2(t)t)g_2(t)t, g_2(t)t^4)}{(1 - p(t))^5} \right] g_2(t), \end{aligned}$$

and

$$\bar{g}_3(t) = g_3(t) - 1.$$

We have that  $\bar{g}_i(0) = -1 < 0$  and  $\bar{g}_i(t) \rightarrow +\infty$  as  $t \rightarrow r_0^-$ . By applying the intermediate value theorem on the functions  $\bar{g}_i$  defined on the interval  $[0, r_0]$ , we deduce that there exist zeros on the interval  $(0, r_0)$ . Denote by  $r_i$  the smallest zero of functions  $\bar{g}_i$  on the interval  $(0, r_0)$ . Define the radius of convergence  $r$  by

$$r = \min\{r_i\}, i = 1, 2, 3. \quad (2.2)$$

Then, for each  $t \in [0, r)$

$$0 \leq g_i(t) < 1, \quad (2.3)$$

and

$$0 \leq p(t) < 1. \quad (2.4)$$

Define also parameter  $R$  by

$$R = \max\{r, r + |b_0|v_0(r)r, g_1(r)r + |b_1|v_0(g_1(r)r)g_1(r)r, g_2(r)r + |b_2|v_0(g_2(r)r)g_2(r)r\}. \quad (2.5)$$

Next, we present the local convergence analysis of method (1.1) under the preceding notation.

**THEOREM 2.1.** Let  $F : D \subset X \rightarrow X$  be a continuous operator. Let also  $[\cdot, \cdot; F] : D \times D \rightarrow L(X)$  be a divided difference of order one. Suppose: there exists  $x^* \in D$  such that operator  $F$  is Fréchet-differentiable at  $x = x^*$ ,

$$F(x^*) = 0, F'(x^*)^{-1} \in L(X); \quad (2.6)$$

there exist functions  $w_0 : [0, +\infty) \times [0, +\infty) \rightarrow [0, +\infty)$ ,  $v_0 : [0, +\infty) \rightarrow [0, +\infty)$  continuous, nondecreasing with  $w_0(0, 0) = 0$  such that for each  $x, y \in D$ ,

$$\|F'(x^*)^{-1}([y, x; F] - F'(x^*))\| \leq w_0(\|y - x^*\|, \|x - x^*\|), \quad (2.7)$$

$$\|[x, x^*; F]\| \leq v_0(\|x - x^*\|); \quad (2.8)$$

there exist functions  $w : [0, r_0) \times [0, r_0) \rightarrow [0, +\infty)$ ,  $v : [0, r_0) \times [0, r_0) \rightarrow [0, +\infty)$  continuous, non-decreasing with  $w(0, 0) = 0$  such that for each  $x, y \in D_0 = D \cap U(x_0, r_0)$

$$\|F'(x^*)^{-1}([x, y; F] - [x, x^*; F])\| \leq w(\|x - x^*\|, \|y - x^*\|) \quad (2.9)$$

$$\|F'(x^*)^{-1}[x, y; F]\| \leq v(\|x - x^*\|, \|y - x^*\|) \quad (2.10)$$

and

$$\overline{U}(x^*, R) \subseteq D \quad (2.11)$$

where  $r_0, r$  and  $R$  are given by (2.1), (2.2) and (2.5) respectively. Then sequence  $\{x_n\}$  generated for  $x_0 \in U(x^*, r) - \{x^*\}$  by method (1.1) is well defined, remains in  $U(x^*, r)$  for each  $n = 0, 1, 2, \dots$  and converges to  $x^*$ . Moreover, the following estimates hold

$$\|y_n - x^*\| \leq g_1(\|x_n - x^*\|)\|x_n - x^*\| \leq \|x_n - x^*\| < r, \quad (2.12)$$

$$\|z_n - x^*\| \leq g_2(\|x_n - x^*\|)\|x_n - x^*\| \leq \|x_n - x^*\|, \quad (2.13)$$

and

$$\|x_{n+1} - x^*\| \leq g_3(\|x_n - x^*\|)\|x_n - x^*\| \leq \|x_n - x^*\|, \quad (2.14)$$

where, the functions  $g_i, i = 1, 2, 3$  are defined previously. Furthermore, if there exists for  $R^* \geq r$ , such that

$$w_0(R^*, 0) < 1 \text{ or } w_0(0, R^*) < 1, \quad (2.15)$$

then, the limit point  $x^*$  is the only solution of equation  $F(x) = 0$  in  $D_1 := D \cap \overline{U}(x^*, R^*)$ .

**Proof.** We shall show estimates (2.12)-(2.14) using induction on  $k = 0, 1, 2 \dots$ . Using (2.6) and (2.8), we have that

$$\begin{aligned} \|u_0 - x^*\| &= \|x_0 - x^* + b_0(F(x_0) - F(x^*))\| \\ &\leq \|x_0 - x^* + b_0[x_0, x^*; F](x_0 - x^*)\| \\ &\leq \|x_0 - x^*\| + |b_0|v_0(\|x_0 - x^*\|)\|x_0 - x^*\| \\ &\leq r + |b_0|v_0(r)r \leq R, \end{aligned} \quad (2.16)$$

so  $u_0 \in \bar{U}(x^*, R)$ . Then, by (2.2), (2.4), (2.5), (2.6), (2.7) and (2.16), we get that

$$\begin{aligned} \|F'(x^*)^{-1}([x_0, u_0; F] - F'(x^*))\| &\leq w_0(\|x_0 - x^*\|, \|u_0 - x^*\|) \\ &\leq w_0(R, R + |b_0|v_0(R)R) \\ &= p(R) < 1. \end{aligned} \quad (2.17)$$

It follows from (2.17) and the Banach Lemma on invertible operators [4-7] that  $A_0 = [x_0, u_0; F]^{-1} \in L(X)$  and

$$\|A_0^{-1}F'(x^*)\| \leq \frac{1}{1 - w_0(\|x_0 - x^*\|, \|u_0 - x^*\|)}. \quad (2.18)$$

We also have that  $y_0, z_0, x_1$  are well defined by the method (1.1) for  $n = 0$ . By (2.2), (2.3)(for  $i = 1$ ), (2.6), (2.9), (2.18) and the first substep of method (1.1) we obtain in turn that

$$\begin{aligned} \|y_0 - x^*\| &= \|x_0 - x^* - A_0^{-1}F'(x_0)\| \\ &\leq \|A_0^{-1}F'(x^*)\| \|F'(x^*)^{-1}(A_0(x_0 - x^*) - F(x_0))\| \\ &\leq \|A_0^{-1}F'(x^*)\| \|F'(x^*)^{-1}(A_0 - [x_0, x^*; F])(x_0 - x^*)\| \\ &\leq \frac{w(\|x_0 - x^*\|, \|x_0 - x^*\| + |b_0|v_0(\|x_0 - x^*\|)\|x_0 - x^*\|)\|x_0 - x^*\|}{1 - p(\|x_0 - x^*\|)} \\ &= g_1(\|x_0 - x^*\|)\|x_0 - x^*\| \leq \|x_0 - x^*\| < r, \end{aligned} \quad (2.19)$$

which shows (2.12) for  $k = 0$  and  $y_0 \in U(x^*, r)$ . As in (2.19) (with  $y_0 = x_0$ ), we obtain the estimates

$$\begin{aligned} &\|y_0 - x^* - A_0^{-1}F(y_0)\| \\ &\leq \|A_0^{-1}F'(x^*)\| \|F'(x^*)^{-1}(A_0 - [y_0, x^*; F])(y_0 - x^*)\| \\ &\leq \frac{w(\|x_0 - x^* + x^* - y_0\|, \|u_0 - x^*\|)\|y_0 - x^*\|}{1 - p(\|x_0 - x^*\|)} \\ &\leq \frac{w(\|x_0 - x^*\| + \|y_0 - x^*\|, \|x_0 - x^*\| + |b_0|v_0(\|x_0 - x^*\|)\|x_0 - x^*\|)\|y_0 - x^*\|}{1 - p(\|x_0 - x^*\|)} \end{aligned} \quad (2.20)$$

and

$$\|(1 - a_0)A_0^{-1}F(y_0)\| \leq \frac{|1 - a_0|v(\|y_0 - x^*\|, 0)\|y_0 - x^*\|}{1 - p(\|x_0 - x^*\|)}. \quad (2.21)$$

In view of the second substep of method (1.1) for  $n = 0$ , (2.2), (2.3) (for  $i = 2$ ), (2.10), (2.19)-(2.21) we get in turn that

$$\begin{aligned}
\|z_0 - x^*\| &\leq \|y_0 - x^* - A_0^{-1}F(y_0)\| + \|(1 - a_0)A_0^{-1}F(y_0)\| + \|(3 - 2a_0)A_0^{-1}B_0A_0^{-1}F(y_0)\| \\
&\quad + \|(a_0 - 2)A_0^{-1}B_0A_0^{-1}F(y_0)\| \\
&\leq \left[ \frac{w(\|x_0 - x^*\| + \|y_0 - x^*\|, \|x_0 - x^*\| + |b_0|v_0(\|x_0 - x^*\|)\|x_0 - x^*\|)}{1 - p(\|x_0 - x^*\|)} \right. \\
&\quad + \frac{|1 - a_0|v(\|y_0 - x^*\|, 0)}{1 - p(\|x_0 - x^*\|)} \\
&\quad + |3 - 2a_0| \frac{v(\|h_0 - x^*\|, \|y_0 - x^*\|)v(\|y_0 - x^*\|, 0)}{(1 - p(\|x_0 - x^*\|))^2} \\
&\quad \left. + |a_0 - 2| \frac{v(\|h_0 - x^*\|, \|y_0 - x^*\|)^2 v(\|y_0 - x^*\|, 0)}{(1 - p(\|x_0 - x^*\|))^3} \right] \|y_0 - x^*\| \\
&\leq g_2(\|x_0 - x^*\|)\|x_0 - x^*\| \leq \|x_0 - x^*\| < r,
\end{aligned} \tag{2.22}$$

which shows (2.13) and  $z_0 \in U(x^*, r)$ . We also used

$$\begin{aligned}
\|h_0 - x^*\| &\leq \|y_0 - x^*\| + |b_1|v_0(\|y_0 - x^*\|)\|y_0 - x^*\| \\
&\leq g_1(r)r + |b_1|v_0(g_1(r)r)g_1(r)r \leq R.
\end{aligned} \tag{2.23}$$

As in (2.20), (2.21) (for  $y_0 = z_0$ ), we obtain the estimates

$$\begin{aligned}
&\|z_0 - x^* - A_0^{-1}F(z_0)\| \\
\leq &\frac{w(\|x_0 - x^*\| + \|z_0 - x^*\|, \|x_0 - x^*\| + |b_0|v_0(\|x_0 - x^*\|, \|x_0 - x^*\|)\|z_0 - x^*\|)}{1 - p(\|x_0 - x^*\|)}.
\end{aligned} \tag{2.24}$$

and

$$\|(1 - a_1)A_0^{-1}F(z_0)\| \leq \frac{|1 - a_1|v(\|z_0 - x^*\|, 0)\|z_0 - x^*\|}{1 - p(\|x_0 - x^*\|)}. \tag{2.25}$$

Then, using the third substep of method (1.1) for  $n = 0$ , (2.2), (2.3) (for  $i = 3$ ), (2.18), (2.19) (2.22), (2.24) and (2.25), we have in turn that

$$\begin{aligned} \|x_1 - x^*\| &\leq \|z_0 - x^* - A_0^{-1}F(z_0)\| + \|(1 - a_1)A_0^{-1}F(z_0)\| \\ &\quad + \|a_2A_0^{-1}C_0A_0^{-1}F(z_0)\| + \|a_3A_0^{-1}C_0A_0^{-1}C_0A_0^{-1}F(z_0)\| \\ &\quad + \|a_4A_0^{-1}C_0A_0^{-1}C_0A_0^{-1}C_0A_0^{-1}(z_0)\| + \|a_5A_0^{-1}C_0A_0^{-1}C_0A_0^{-1}C_0A_0^{-1}C_0A_0^{-1}F(z_0)\| \\ &\leq \left[ \frac{w(\|x_0 - x^*\| + \|z_0 - x^*\|, \|x_0 - x^*\| + |b_0|v_0(\|x_0 - x^*\|)\|x_0 - x^*\|)}{1 - p(\|x_0 - x^*\|)} \right. \\ &\quad + \frac{|1 - a_1|v(\|z_0 - x^*\|, 0)}{1 - p(\|x_0 - x^*\|)} \\ &\quad + |a_2| \frac{v(\|z_0 - x^*\| + |b_2|v_0(\|z_0 - x^*\|)\|z_0 - x^*\|, \|z_0 - x^*\|)}{(1 - p(\|x_0 - x^*\|))^2} \\ &\quad + |a_3| \frac{v(\|z_0 - x^*\| + |b_2|v_0(\|z_0 - x^*\|)\|z_0 - x^*\|, \|z_0 - x^*\|)^2}{(1 - p(\|x_0 - x^*\|))^3} \\ &\quad + |a_4| \frac{v(\|z_0 - x^*\| + |b_2|v_0(\|z_0 - x^*\|)\|z_0 - x^*\|, \|z_0 - x^*\|)^3}{(1 - p(\|x_0 - x^*\|))^4} \\ &\quad \left. + |a_5| \frac{v(\|z_0 - x^*\| + |b_2|v_0(\|z_0 - x^*\|)\|z_0 - x^*\|, \|z_0 - x^*\|)^4}{(1 - p(\|x_0 - x^*\|))^5} \right] \\ &\leq g_3(\|x_0 - x^*\|)\|x_0 - x^*\| \leq \|x_0 - x^*\| < r, \end{aligned} \tag{2.26}$$

which shows (2.14) for  $n = 0$  and  $x_1 \in U(x^*, r)$ . We also used that

$$\begin{aligned} \|l_0 - x^*\| &\leq \|z_0 - x^*\| + |b_2|v_0(\|z_0 - x^*\|)\|z_0 - x^*\| \\ &\leq g_2(r)r + |b_2|v_0(g_2(r)r)g_2(r)r \leq R. \end{aligned} \tag{2.27}$$

The induction for error bounds (2.12)-(2.14) is completed in an analogous way, if we replace  $x_0, y_0, z_0, x_1$  by  $x_k, y_k, z_k, x_{k+1}$  in the preceding estimates. Then, in view of the estimate

$$\|x_{k+1} - x^*\| \leq c\|x_k - x^*\| < r, \tag{2.28}$$

here  $c = g_3(\|x_0 - x^*\|) \in [0, 1)$ , we deduce that  $\lim_{k \rightarrow \infty} x_k = x^*$  and  $x_{k+1} \in U(x^*, r)$ . Finally, to show the uniqueness of the solution  $x^*$  in  $D_1$ , suppose  $y^* \in D_1$  with  $F(y^*) = 0$ . Define the linear operator  $Q = [y^*, x^*; F]$ . Using (2.7) and (2.15) we obtain that

$$\|F'(x^*)^{-1}(Q - F'(x^*))\| \leq w_0(\|y^* - x^*\|, 0) \leq w_0(R^*, 0) < 1, \tag{2.29}$$

so  $Q^{-1} \in L(X)$ . Then, using the identity

$$0 = F(y^*) - F(x^*) = Q(y^* - x^*), \tag{2.30}$$

we conclude that  $x^* = y^*$ . □

**REMARK 2.2.** (a) A stronger condition depending on function  $w_0$  can replace condition (2.8). Indeed, using (2.7), we have in turn that

$$\begin{aligned} \|[x, x^*; F]\| &= \|F'(x^*)F'(x^*)^{-1}[x, x^*; F]\| \\ &\leq \|F'(x^*)\| \|[F'(x^*)^{-1}([x, x^*; F] - F'(x^*)) + I]\| \\ &\leq \beta(1 + \|F'(x^*)^{-1}([x, x^*; F] - F'(x^*))\|) \\ &\leq \beta(1 + w_0(t, 0)). \end{aligned}$$

Therefore,  $v_0(t) := \beta(1 + w_0(t, 0))$  or  $v_0(t) := \beta(1 + w_0(r_0, 0))$  or  $v_0(t) := \beta(1 + w_0(r_0, 0))$ , where  $\|F'(x^*)\| \leq \beta$ .

(b) It is worth noticing that method (1.1) are not changing if we use the new instead of the old conditions [2]. Moreover, for the error bounds in practice we can use the computational order of convergence (COC) [25]

$$\xi = \frac{\ln \frac{\|x_{n+2} - x^*\|}{\|x_{n+1} - x^*\|}}{\ln \frac{\|x_{n+1} - x^*\|}{\|x_n - x^*\|}}, \quad \text{for each } n = 1, 2, \dots$$

or the approximate computational order of convergence (ACOC)

$$\xi^* = \frac{\ln \frac{\|x_{n+2} - x_{n+1}\|}{\|x_{n+1} - x_n\|}}{\ln \frac{\|x_{n+1} - x_n\|}{\|x_n - x_{n-1}\|}}, \quad \text{for each } n = 0, 1, 2, \dots$$

instead of the error bounds obtained in Theorem 2.1.

(c) As in [2] we can consider methods with memory, if we approximate  $b_0 = b_1 = b_2 := T^{(n)} = -[u_{n-1}, x_{n-1}; F]^{-1}$ ,  $n \geq 1$ . This way the order of convergence  $4 + \sqrt{19} \approx 8.3589$  was shown in [2, Theorem 2]. The method with memory is defined by  $B_0^{(k)} = B_1^{(k)} = B_2^{(k)} = -[u_{n-1}, x_{n-1}; F]^{-1} = -A_{n-1}^{-1}$

$$\begin{aligned} y_n &= x_n - q_1 \\ z_n &= y_n - 3q_2 + 3q_3 - q_4 \\ x_{n+1} &= z_n - 4q_5 + 6q_6 - 4q_7 + q_8, \end{aligned} \quad (2.31)$$

where

$$u_n = x_n + B_0^{(n)}F(x_n), \quad h_n = y_n + B_1^{(n)}F(y_n), \quad l_n = z_n + B_2^{(n)}F(z_n).$$

In order for us to apply Theorem 2.1 to method (2.31), let us simply replace  $b_0, b_1, b_2$  in Theorem 2.1 by (see (2.7))

$$\frac{\beta_0}{1 - p(\|x_0 - x^*\|)} \leq \frac{\beta_0}{1 - p(r)} := b(r) \quad (2.32)$$

where  $\|F'(x^*)^{-1}\| \leq \beta_0$ . Indeed, for example by (2.7)

$$|b_2| \leq \| -[u_{n-1}, x_{n-1}; F]^{-1}F'(x^*)F'(x^*)^{-1} \|$$

$$\|[u_{n-1}, x_{n-1}; F]^{-1}F'(x^*)\| \|F'(x^*)^{-1}\| \leq \frac{\beta_0}{1 - p(\|x_0 - x^*\|)} \leq \frac{\beta_0}{1 - p(r)} = b(r). \quad (2.33)$$

The other methods studied in [2] can also be handled in an analogous way as method (2.31).

### 3. NUMERICAL EXAMPLES

The numerical examples are presented in this section. We choose

$$[x, y; F] = \int_0^1 F'(y + \theta(x - y))d\theta.$$

We have taken  $b_0 = 0.5, a_1 = 4, a_2 = -6; a_3 = 4, a_4 = -1$  and  $a_5 = 0$ .



**EXAMPLE 3.1.** Let  $X = \mathbb{R}^3$ ,  $D = \bar{U}(0, 1)$ ,  $x^* = (0, 0, 0)^T$ . Define function  $F$  on  $D$  for  $w = (x, y, z)^T$  by

$$F(w) = (e^x - 1, \frac{e-1}{2}y^2 + y, z)^T.$$

Then, the Fréchet-derivative is given by

$$F'(v) = \begin{bmatrix} e^x & 0 & 0 \\ 0 & (e-1)y + 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}.$$

Notice that using the (2.9) conditions, we get  $w_0(s, t) = \frac{L_0}{2}(s+t)$ ,  $v(t) = \frac{1}{2}(1+e)$ ,  $w(s, t) = \frac{L_0}{2}t$ ,  $v(s, t) = \frac{1}{2}e^{\frac{1}{L_0}(s+t)}$ ,  $L_0 = e - 1$ . The parameters are

$$r_1 = 0.3394, r_2 = 0.3576, r_3 = 0.3086 = r.$$

**EXAMPLE 3.2.** Returning back to the motivational example given at the introduction of this study, we can choose (see also Remark 2.2 (a) for function  $v$ )  $w_0(t, s) = w(t, s) = \frac{t+s}{16}$  and  $v_0(t) = v(s, t) = 1 + w_0(0, t)$  and  $\beta = 1$ . Then, the radius of convergence  $r$  is given by

$$r_1 = 3.081318457, r_2 = 0.99469429615, r_3 = 0.1667265158 = r.$$

## REFERENCES

- [1] M. F. Abad, A. Cordero, J. R. Torregrosa, Fourth and Fifth-order methods for solving nonlinear systems of equations: An application to the global positioning system *Abstract and Applied Analysis*, (2013),
- [2] F. Ahmad, F. Soleymani, F. K. Haghani, S.S. Capizzano, Higher order derivative-free iterative methods with and without memory for systems of nonlinear equations, *Appl. Math. Comput.*, —
- [3] S. Amat, M.A. Hernández, N. Romero, Semilocal convergence of a sixth order iterative method for quadratic equations, *Applied Numerical Mathematics*, 62 (2012), 833-841.
- [4] I.K. Argyros, Computational theory of iterative methods. Series: Studies in Computational Mathematics, 15, Editors: C.K.Chui and L. Wuytack, Elsevier Publ. Co. New York, U.S.A, 2007.
- [5] Argyros, I.K., A semilocal convergence analysis for directional Newton methods. *Math. Comput.* 80 (2011), 327–343.
- [6] I.K. Argyros, S. Hilout, Weaker conditions for the convergence of Newton's method. *J. Complexity* 28 (2012) 364–387.
- [7] I. K. Argyros and Said Hilout, Computational methods in nonlinear analysis. Efficient algorithms, fixed point theory and applications, World Scientific, 2013.
- [8] I.K. Argyros and H. Ren, Improved local analysis for certain class of iterative methods with cubic convergence, *Numerical Algorithms*, 59(2012), 505-521.
- [9] A. Cordero, F. Martinez, J. R. Torregrosa, Iterative methods of order four and five for systems of nonlinear equations, *J. Comput. Appl. Math.* 231, (2009), 541-551.
- [10] A. Cordero, J. Hueso, E. Martinez, J. R. Torregrosa, A modified Newton-Tarratt's composition, *Numer. Algor.* 55, (2010), 87-99.
- [11] A. Cordero, J. R. Torregrosa, M. P. Vasileva, Increasing the order of convergence of iterative schemes for solving nonlinear systems, *J. Comput. Appl. Math.* 252, (2013), 86-94.

- [12] G.M Grau-Sanchez, A.Grau, M .Noguera, On the computational efficiency index and some iterative methods for solving systems of non-linear equations, *J. Comput. Appl Math.*, 236 (2011), 1259-1266.
- [13] J. M. Gutiérrez, A.A. Magreñán and N. Romero, On the semi-local convergence of Newton-Kantorovich method under center-Lipschitz conditions, *Applied Mathematics and Computation*, 221 (2013), 79-88.
- [14] L.V. Kantorovich, G.P. Akilov, *Functional Analysis*, Pergamon Press, Oxford, 1982.
- [15] J.S. Kou, Y. T. Li and X.H. Wang, A modification of Newton method with third-order convergence, *Appl. Math. Comput.* 181, (2006), 1106-1111.
- [16] A. A. Magrenan, Different anomalies in a Jarratt family of iterative root finding methods, *Appl. Math. Comput.* 233, (2014), 29-38.
- [17] A. A. Magrenan, A new tool to study real dynamics: The convergence plane, *Appl. Math. Comput.* 248, (2014), 29-38.
- [18] M. S. Petkovic, B. Neta, L. Petkovic, J. Džunič, *Multipoint methods for solving nonlinear equations*, Elsevier, 2013.
- [19] F. A. Potra and V. Pták, *Nondiscrete Induction and Iterative Processes*, in: *Research Notes in Mathematics*, Vol. 103, Pitman, Boston, 1984.
- [20] A.N .Romero, J.A. Ezquerro, M .A. Hernandez, *Approximacion de soluciones de algunas ecuaciones integrals de Hammerstein mediante metodos iterativos tipo. Newton*, XXI Congresode ecuaciones diferenciales y aplicaciones Universidad de Castilla-La Mancha (2009)
- [21] W.C. Rheinboldt, An adaptive continuation process for solving systems of nonlinear equations, In: *Mathematical models and numerical methods* (A.N.Tikhonov et al. eds.) pub.3, (1977), 129-142 Banach Center, Warsaw Poland.
- [22] J.R. Sharma, P.K. Guha and R. Sharma, An efficient fourth order weighted-Newton method for systems of nonlinear equations, *Numerical Algorithms*, 62, 2, (2013), 307-323.
- [23] J.R. Sharma, P.K. Guha, An efficient fifth order method for systems of nonlinear equations, *Comput. Math. Appl* 67(2014), 591-601.
- [24] J.F.Traub, *Iterative methods for the solution of equations*, AMS Chelsea Publishing, 1982.
- [25] S. Weerakoon, T.G.I. Fernando, A variant of Newton's method with accelerated third-order convergence, *Appl. Math. Lett.* 13, (2000), 87-93.